

Recommended protocol for CK20

Obtained in run 62

25 Feb 2021

Immunostainer

Type: Manual

Primary antibody

Clone: ZM42
Producer: Zeta Corporation
Product no. / lot no.: Z2349 / 1806083
Diluent: Normal Antibody Diluent
Diluent producer / no.: Zeta Corporation / ZD19
Dilution factor: 1:200
Incubation time / temperature: 45 min. / 25°C

Epitope retrieval, HIER

Device: Pressure Cooker
Buffer: Zeta Corporation Tris-EDTA HIER Solution pH 9.0
Heating time at max. temp.: 45 min.
Maximum heating temp.: 95°C

Visualization system

Producer: Zeta Corporation
Product / no: Zeta Universal Polymer / ZD8
Linker: None
Incubation time polymer: 30 min.
Incubation temperature: 25°C

Chromogen

Producer: Zeta Corporation
Product / no: DAB Chromogen / ZD11
Incubation time / temperature: 5 min. / 25°C
Enhancement: None

Disclaimer:

NordiQC makes every attempt to provide accurate and up-to-date information, yet NordiQC does not make any claim or warranty regarding the accuracy of the provided information nor does it represent that the contents of the web site and protocols reflect the most recent developments in immunohistochemistry at any point in time.